

This chapter appeared in Easton, A, & Emery, N (eds) (2005)

The Cognitive Neuroscience of Social Behaviour, Psychology Press.

Autism and the origins of social neuroscience.

Simon Baron-Cohen

Autism Research Centre
Departments of Psychiatry and Experimental Psychology,
University of Cambridge,
Douglas House, 18b Trumpington Rd,
Cambridge, CB2 2AH UK

Acknowledgments: The MRC, the National Alliance for Autism Research, Cure Autism Now, and the James S McDonnell Foundation provided valuable support during the period of this work. Portions of this chapter appeared in Baron-Cohen et al (2002).

‘Social neuroscience’ is something of a new phrase, and the editors of this volume are to be congratulated for collecting together the fragmented work, and thereby helping the creation of a new field. In their introduction to this book, they cover the history of this idea; but for me there are some themes – even lessons - worth highlighting.

Lessons from history

Non-social neuroscience

When I started in psychology some twenty years ago, there was almost no hint of social neuroscience. Cognitive neuroscience was alive and well, so this was not a reflection of a lack of activity in the wider field. Rather, it reflects that studies of the brain were for the most part *non-social*. We had, as examples, Blakemore’s and Weizcrantz’ classic studies of the visual system in kittens, monkeys, and humans to tell us which (non-social) features of the environment were perceived and how. We had Luria and Shallice’s classic studies of the (non-social) control of action to reveal not just a ‘central executive’ for planning in the brain, but a syndrome of executive dysfunction. We had a wealth of other studies investigating conditions such as amnesia and agnosia to tell us how memory and knowledge of information *in general* worked in the brain. Even Wernicke’s and Broca’s classic studies of the language system in brain-damaged patients focused for the most part on the production and comprehension of words *in general* by the normal brain. But such aphasias were

lexical or syntactic or semantic, but ignored the social aspects of communication: pragmatics.

Why was this? After all, cognitive neuroscientists then weren't fools. They knew then, just we know now, that the human brain, and indeed most primate brains, exist *first and foremost* in a social world. Primates do not sit in solitary, solipsistic universes. So why did they treat the brain as if it had no special interest in the social world?

My guess is that there are (at least) two explanations for this. First, cognitive neuroscience followed a parsimonious approach of assuming that the brain is a *general* information processor. Whether tacitly or explicitly, the assumption has been that the visual system, or the auditory system, or the memory systems, or the planning system, work on input of a general kind, where content plays little role. Of course, distinctions have been drawn, such as visual vs. auditory memory, but within a given system it was held to not matter whether the visual input is a tree or a car: the search was to identify the general operating principles of the visual system. The same applies to memory. It matters little if we are studying memory of cars or of animals. We should still be able to identify the general operating principles of the memory system.

Such a content-free approach was laudable in its parsimony, as the danger otherwise was that neuroscience could have ended up positing a very large number of specialized circuits for different classes of information, and then the whole enterprise of understanding the basic laws of the brain would have been thrown off course. However, throughout this enterprise, there were always cracks appearing in this 'brain-as-a-general-processor' theory. Just one example was the case of prosopagnosia,

where some clinicians claimed that some patients could recognize any kind of object *except* faces. And the publication of Fodor's landmark book on *modularity* still stands as a major challenge to such a general theory (Fodor, 1983).

The second possible explanation relates to cognitive neuroscientists as natural scientists. The nature of natural science is to try to isolate variables in a system under controlled conditions. The ultimate model for natural scientists is physics, and it is no surprise that even in the study of the human brain, the dominant approach has been to study how the brain responds to the manipulation of elementary features of the input. Is a vertical edge detected by the same assembly of neurons as a horizontal edge? Is a regular verb processed in the same way as an irregular verb? Again, such a focus on controllable, simple stimuli or features is laudable, since in this way one can make inferences about how the system works. If one were dealing with the complexity of the social world, how on earth could one begin to isolate what was causing what?

This is not intended as a criticism of cognitive neuroscience in adopting a general, or a non-social approach. The natural science methodology has reaped great benefits and has re-appropriated the study of the mind from the hands of psychoanalysts and social scientists who ignored the brain and biology for decades. We have much to be grateful for. But there may be a set of parallels that emerge from a range of fields within psychology that show a similar disregard of the special nature of the social world. The lessons have been learned late.

Non-social psychology

The study of child development began with a disproportionate focus on the non-social aspects of cognition. Piaget's classic studies of object permanence and what is really 'folk physics' dominated until the 1980s, until Bruner (in Oxford) and his students reminded the field that children have minds that are trying to make sense of a social world, and not just a physical world. Indeed, the shift of focus to the pragmatics of communication (and away from traditional, Chomskyan approaches to language acquisition), and the 'discovery' of the developing child's 'theory of mind' (Astington, Harris, & Olson, 1988; Wellman, 1990), owes a lot to Bruner's repeated concern that we were treating the child-as-scientist and ignoring the child-in-relationships (Bruner, 1983).

The same history unfolded in the field of intelligence. Almost all the early and classical IQ tests sought to assess the person's non-social IQ: David Wechsler's non-verbal subtests of object-assembly or block design, or his verbal subtests of vocabulary or digit span, or Raven's matrices, taught us an enormous amount about the predictive power of IQ (Raven, 1956; Wechsler, 1939), but virtually ignored what today is called "social intelligence" or "emotional intelligence" (Goleman, 1995) Equally, cognitive psychology focused in large part on the non-social aspects of cognition, with the new field of "social cognition" only coming in quite late in the 20th century (Shantz, 1983).

If we look at the field of primatology we can see a similar pattern. The attempt to understand the evolution of the intelligence and the evolution of the brain focused on humans-as-tool-users and general problem-solvers (or early hominids as 'folk

physicists') as the driving force behind the evolution of a larger, more powerful brain . It was rather late in the 20th century when it was asked "Does the chimpanzee have a theory of mind?" (Premack & Woodruff, 1978), and that it was proposed that the driving force behind the evolution of intelligence and the brain may have been the need to socially outwit competitors (the "machiavellian intelligence hypothesis" (Whiten, 1997)).

I lay out this brief and partial history because I think may be lessons to be learnt. Naturally, there is a risk of painting the history as too black and white. Too non-social when all along there was a streak of social neuroscience running through it. We know that Piaget's concept of 'egocentrism' applied not only to the child's folk physics (to explain the child's errors in understanding conservation of mass, for example) but also to the child's folk psychology (to explain the child's errors in communication). And we know that Charles Gross's classic single-cell recording studies were not only identifying cell-assemblies that fired in response to non-social aspects of the visual environment, but also in response to specifically social features such as hands and faces (Gross, Rocha-Miranda, & Bender, 1972). We know that Harry Harlow's, Robert Hinde's and John Bowlby's classic studies of the attachment system in monkeys and humans progressed despite this history, and indeed ethology never lost sight of its social context (Bowlby, 1969). But these exceptions to the rule do not, I think, invalidate the broad picture I have painted. Rather, they were the seeds for the new field of social neuroscience.

From non-social to social accounts of autism

The study of autism has followed a similar history. The psychological theories of autism before the 1980s were for the most part non-social. The child's social difficulties were attributed to either a failure to generalize (Rimland, 1964) or were seen as secondary to a language disorder (Rutter, 1978), or were thought to reflect a failure to process meaning (or semantics) (Hermelin & O'Connor, 1970), for example. For this reason, the proposal (by my colleagues and I in the 1980s) that the social and communication difficulties that are the hallmark of autism might reflect a *specific* deficit in an aspect of *social cognition* was treated as quite novel. Our idea was that there might be specific brain regions or neural circuits that underpinned social understanding, and ultimately social behaviour. We opened this area of investigation by asking "Does the autistic child have a theory of mind?" (Baron-Cohen, Leslie, & Frith, 1985). A related investigation into emotion recognition in autism (Hobson, 1986) was also regarded as new and important. Later in this chapter I summarise how this work has unfolded in the subsequent 20 years.

The Social Brain

My own theoretical and empirical work was greatly enriched by Leslie Brothers' important proposal of a network of neural regions that comprise the "social brain" (Brothers, 1990). She suggested this included areas of prefrontal cortex (orbital and medial areas particularly), the superior temporal sulcus, and the amygdala. Since the neuro-developmental condition of autism involves deficits in what today I refer to as 'empathizing', (Baron-Cohen, 2002), it is plausible that autism may be caused by an abnormality in one or more of these brain areas.

The idea that social understanding might be independent of general intelligence comes from 3 sources:

- There are individuals who are capable of considerable understanding of the non-social world (e.g., physics, maths, engineering) yet who readily admit to finding the social world confusing (Baron-Cohen, Wheelwright, Stone, & Rutherford, 1999; Sacks, 1994)
- The opposite type of individual also exists: people who have no difficulty interacting with the social world but who find non-social problem-solving confusing (Karmiloff-Smith, Grant, Bellugi, & Baron-Cohen, 1995).
- Certain kinds of brain damage (e.g., to the amygdala) can cause selective impairment in social judgment (Damasio, Tranel, & Damasio, 1990) without any necessary loss to general problem-solving ability. Loss of social judgment can of course co-occur with memory and executive dysfunction (Tranel & Hyman, 1990), but the functional double dissociation between empathizing and non-social aspects of intelligence suggests their neural independence.

In the remainder of this chapter I review the evidence for the normal development of empathizing. I then review the literature suggesting autism involves an empathizing deficit. Finally, I end with a summary of the evidence for the role of the amygdala in empathy. The evidence for the social function of orbito- and medial prefrontal cortex, and the superior temporal sulcus, is reviewed elsewhere (Baron-Cohen & Ring, 1994; Baron-Cohen et al., 1994).

The empathizing theory of autism

Autism is diagnosed when a child or adult has abnormalities in a 'triad' of behavioural domains: social development, communication, and repetitive behaviour/obsessive interests (A.P.A, 1994; I.C.D-10, 1994) . Asperger Syndrome (AS) was first described by Asperger (Asperger, 1944). The descriptions of the children he documented overlapped considerably with the accounts of childhood autism (Kanner, 1943). Little was published on AS in English until relatively recently (Frith, 1991; Wing, 1981). Current diagnostic practice recognises people with AS as meeting the same criteria as for high-functioning autism (HFA), but with no history of language delay, and with no cognitive delay.

The mindblindness theory of autism (Baron-Cohen, 1995), and its extension into empathizing theory (Baron-Cohen, 2002) proposes that in autism spectrum conditions there are deficits in the normal process of empathizing, relative to mental age. These deficits can occur by degrees. The term 'empathizing' encompasses the following earlier terms: 'theory of mind', 'mind-reading', and taking the 'intentional stance' (Dennett, 1987).

Empathizing involves two major elements: (a) the ability to attribute mental states to oneself and others, as a natural way to understand agents (Baron-Cohen, 1994; Leslie, 1995; Premack, 1990); (b) having an emotional reaction that is appropriate to the other person's mental state. In this sense, it includes what is normally meant by the

term 'theory of mind' (the attributional component) but it goes beyond this, to also include having some affective reaction (such as sympathy).

The first of these, the mental state attribution component, this has been widely discussed in terms of being an evolved ability, the response of a cognitive system to a universe that can be broadly divided into two kinds of entities: those that do and do not possess intentionality (Brentano, 1970). The mental state attribution component is effectively judging if this is the sort of entity that might possess intentionality. Intentionality is defined as the capacity of something to refer or point to things other than itself. A chair cannot point to anything. It just is. In contrast, a rabbit can 'look' at a carrot, it can 'want' the carrot, and it can 'think' that this is a carrot, etc. Essentially, agents have intentionality, whereas non-agents do not. This means that when we observe agents and non-agents move, we construe their motion as having different causes (Csibra, Gergely, Biro, Koos, & Brockbanck, 1999; Gelman & Hirschfield, 1994). Agents can move by self-propulsion, which we naturally interpret as driven by their goals and desires, whilst non-agents cannot.

The second of these, the affective reaction component, is closer to what we ordinarily refer to with the English word 'empathy'. Thus, we not only attribute a mental state to the agent in front of us (e.g., the man is in pain), but we also react to his emotional state with an appropriate emotion ourselves (we feel sorry for him). Empathizing thus essentially allows us to make sense of the behaviour of another agent we are observing, predict what they might do next, and how they might feel. And it allows us to feel connected to another agent's experience, and respond appropriately to them.

The normal development of empathizing

Empathizing develops from human infancy (Johnson, 2000). In the infancy period, it includes

- being able to judge if something is an agent or not (Premack, 1990);
- being able to judge if another agent is looking at you or not (Baron-Cohen & Goodhart, 1994);
- being able to judge if an agent is expressing a basic emotion (Ekman, 1992), and if so, what type.
- engaging in shared attention, for example by following gaze or pointing gestures (Mundy & Crowson, 1997; Scaife & Bruner, 1975; Tomasello, 1988);
- showing concern or basic empathy at another's distress, or responding appropriately to another's basic emotional state (Yirmiya, Sigman, Kasari, & Mundy, 1992);
- being able to judge an agent's goal or basic intention (Premack, 1990).

Empathizing can be identified and studied from at least 12 months of age (Baron-Cohen, 1994; Premack, 1990). Thus, infants dishabituate to actions of 'agents' who appear to violate goal-directedness (Gergely, Nadasdy, Gergely, & Biro, 1995; Rochat, Morgan, & Carpenter, 1997). They also expect agents to 'emote' (express emotion), and expect this to be consistent across modalities (between face and voice) (Walker, 1982). They are also highly sensitive to where another person is looking, and by 14 months will strive to establish joint attention (Butterworth, 1991; Hood, Willen,

& Driver, 1997; Scaife & Bruner, 1975). By 14 months they also start to produce and understand pretence (Bates, Benigni, Bretherton, Camaioni, & Volterra, 1979; Leslie, 1987). By 18 months they begin to show concern at the distress of others (Yirmiya et al., 1992). By 2 years old they begin to use mental state words in their speech (Wellman & Bartsch, 1988).

Empathizing of course develops beyond early childhood, and continues to develop throughout the lifespan. These later developments include:

- attribution of the range of mental states to oneself and others, including pretence, deception, belief (Leslie & Keeble, 1987).
- recognizing and responding appropriately to complex emotions, not just basic ones (Harris, Johnson, Hutton, Andrews, & Cooke, 1989).
- linking mental states to action, including language, and therefore understanding and producing pragmatically appropriate language (Tager-Flusberg, 1993)
- making sense of others' behaviour, and predicting it, and even manipulating it (Whiten, 1991).
- judging what is appropriate in different social contexts, based on what others will think of our own behaviour.
- communicating an empathic understanding of another mind.

Thus, by 3 years old, children can understand relationships between mental states such as seeing leads to knowing (Pratt & Bryant, 1990). By 4 years old they can understand that people can hold false beliefs (Wimmer & Perner, 1983). By 5-6 years old they can

understand that people can hold beliefs about beliefs (Perner & Wimmer, 1985). By 7 years old they begin to understand what not to say in order to avoid offending others (Baron-Cohen, O'riordan, Jones, Stone, & Plaisted, 1999). With age, mental state attribution becomes increasingly more complex (Baron-Cohen, Jolliffe, Mortimore, & Robertson, 1997; Happe, 1993). The little cross-cultural evidence that exists suggests a similar picture in very different cultures (Avis & Harris, 1991).

These developmental data have been interpreted in terms of an innate module being part of the infant cognitive architecture. This has been dubbed a theory of mind mechanism (ToMM) (Leslie, 1995). But as we have suggested, empathizing also encompasses the skills that are involved in normal reciprocal social relationships (including intimate ones) and in sensitive communication. Empathizing is a narrowly defined domain, namely, *understanding and responding to people's minds*. Deficits in empathizing are referred to as degrees of mindblindness.

Empathizing in autism spectrum conditions

Since the first test of mindblindness in children with autism (Baron-Cohen et al., 1985), there have been more than 30 experimental tests. The vast majority of these have revealed profound impairments in the development of their empathizing ability. These are reviewed elsewhere (Baron-Cohen, 1995; Baron-Cohen, Tager-Flusberg, & Cohen, 1993) but include deficits in the following:

- joint attention (Baron-Cohen, 1989a);
- use of mental state terms in language (Tager-Flusberg, 1993);

- production and comprehension of pretence (Baron-Cohen, 1987; Wing & Gould, 1979);
- understanding that “seeing-leads-to-knowing” (Baron-Cohen & Goodhart, 1994; Leslie & Frith, 1988);
- distinguishing mental from physical entities (Baron-Cohen, 1989; Ozonoff, Pennington, & Rogers, 1990);
- making the appearance-reality distinction (Baron-Cohen, 1989b);
- understanding false belief (Baron-Cohen et al., 1985);
- understanding beliefs about beliefs (Baron-Cohen, 1989c);
- understanding complex emotions (Baron-Cohen, 1991);
- showing concern at another’s pain (Yirmiya et al., 1992).

Some children and adults with AS only show their empathizing deficits on age-appropriate adult tests (Baron-Cohen, Jolliffe et al., 1997; Baron-Cohen, Wheelwright, Hill, Raste, & Plumb, 2001; Baron-Cohen, Wheelwright, & Jolliffe, 1997), or on age-appropriate screening instruments such as the Empathy Quotient (EQ) (Baron-Cohen, Richler, Bisarya, Gurunathan, & Wheelwright, 2003; Baron-Cohen & Wheelwright, 2004).

Evidence for the contribution of the amygdala in the social brain and in autism

There are several important lines of evidence implicating the amygdala in primate social behaviour. Extensive reviews exist elsewhere (Kling & Brothers, 1992). We also know that the human amygdala is activated in humans when decoding signals of social importance, such as gaze, expression-recognition (especially of fearful faces), and body movements) (Baron-Cohen, Ring et al., 1999; Bonda, Petrides, Ostry, & Evans, 1996; Kawashima et al., 1999; Morris et al., 1996; Whalen et al., 1998; Wicker, Michel, Henaff, & Decety, 1998). But there are 6 lines of evidence for an amygdala deficit in autism.

(a) Post-mortem evidence A neuroanatomical study of autism at post-mortem found microscopic pathology (in the form of increased cell density) in the amygdala, in the presence of normal amygdala volume (Bauman & Kemper, 1994; Rapin & Katzman, 1998).

(b) An animal model of autism The only animal model of autism involves ablation of the amygdala (in rhesus monkeys) (Bachevalier, 1991). There are obviously limits to any animal model of autism, given that the syndrome involves deficits in higher-order cognition, but Bachevalier makes the case that the effects of amygdala lesions in monkeys resemble some of the symptoms of autism. In particular, the Kluver-Bucy syndrome seems a fairly good animal model of autism (Hetzler & Griffin, 1981).

(c) Similarities between autism and patients following amygdalotomy Patients with amygdala lesions show impairments in social judgement (Adolphs, Tranel, Damasio, &

Damasio, 1994; Young, Hellawell, De Wal, & Johnson, 1996) that have been likened to “acquired autism” (Stone, 2000). The age of onset of deficits in acquired vs. idiopathic cases is likely to mean that the two syndromes also differ in many ways, too. Similarly, patients with autism tend to show a similar pattern of deficits to those seen in patients with amygdala lesions (Adolphs, Sears, & Piven, 2001).

(d) The effects of temporal lobe tubers In cases of tuberous sclerosis, autistic comorbidity is determined by hamartomata in the temporal lobe (Bolton & Griffiths, 1997)¹.

(e) Structural neuroimaging A structural magnetic resonance imaging study of autism reported reduced amygdala volume (Abell et al., 1999).

(f) Functional neuroimaging Using single photon emission computed tomography (SPECT), patients with autism spectrum conditions show significant reductions in temporal lobe blood flow. This is not simply an effect of temporal lobe epilepsy (Gillberg, Bjure, Uvebrant, Vestergren, & Gillberg, 1993). In our earlier functional magnetic resonance imaging (fMRI) study, we found that adults with high functioning autism (HFA) or Asperger Syndrome (AS) showed significantly less amygdala activation during an empathizing task (the ‘Reading the mind in the Eyes’ task), compared to normal controls (Baron-Cohen, Ring et al., 1999) (Baron-Cohen et al., 1999). Adults with HFA or AS, with intelligence in the normal range, show deficits on

¹ We emphasize the amygdala theory of autism, though some of the lines of evidence cited here implicate temporal lobe structures, which include the amygdala but also include other adjacent mesiotemporal areas. It remains for future work to establish the specificity of an amygdala deficit in autism.

this task (Baron-Cohen, Jolliffe et al., 1997; Baron-Cohen et al., 2001), as do parents of children with autism/AS (Baron-Cohen & Hammer, 1997). Children with William's Syndrome are not impaired on this test, despite their general retardation (Tager-Flusberg, Boshart, & Baron-Cohen, 1998).

Other brain areas that might be abnormal in autism

Whilst this chapter highlights the role an amygdala abnormality plays in autism, we do not suggest that this is the only abnormal neural region. For example, the case has been made for anomalous functioning in the cerebellum (Courchesne et al., 1994), hippocampal formation (De Long, 1992), medial frontal cortex (Happe et al., 1996), and fronto-limbic connections (Bishop, 1993) in autism. Reduced neuron size and increased cell-packing density has also been found in the limbic system, specifically the hippocampus, subiculum, entorhinal cortex, amygdala, mammillary bodies, anterior cingulate, and septum in autism (Bauman & Kempner, 1994; Bauman & Kempner, 1985; Bauman & Kempner, 1988; Bauman & Kempner, 1986; Raymond, Bauman, & Kempner, 1996). A full review of neuroimaging of autism may be found elsewhere (Filipek, 1999). Here, we instead follow a line of argument begun by other authors emphasising an amygdala theory of autism (Bachevalier, 1994; Baron-Cohen et al., 2000; Bauman & Kempner, 1988; Hetzler & Griffin, 1981). In the closing section of this chapter we briefly turn from brain regions to the neurochemistry, and particularly the neuroendocrinology of social development.

Foetal testosterone and brain development

Foetal testosterone (FT) acts on the developing brain to influence cerebral lateralisation (Kimura, 1999; Wilson, Foster, Kronenberg, & Larsen, 1998). Evidence for this derives from both animal studies (Arnold & Gorski, 1984; Harris & Levine, 1962; Williams, Barnett, & Meck, 1990), and the effects of abnormal hormonal environments during human pregnancy, such as Congenital Adrenal Hyperplasia or synthetic hormone injections (Collaer & Hines, 1995; Hines & Shipley, 1984).

There is reason to believe that sex hormones might be inversely related to social and language development (Geschwind & Galaburda, 1987; Geschwind & Galaburda, 1985; Geschwind & Galaburda, 1985; Geschwind & Galaburda, 1985). Sex differences (female superiority) have been found in studies of normal language and social development (Baron-Cohen, 2002; Connellan, Baron-Cohen, Wheelwright, Ba'tki, & Ahluwalia, 2001; Hyde & Linn, 1988; Maccoby & Jacklin, 1974), and recent studies suggest an inverse correlation between levels of foetal testosterone as measured in amniotic fluid, with both amount of eye contact measured at 12 months old (Lutchmaya, Baron-Cohen, & Raggatt, 2002a) and vocabulary size at 18 and 24 months old (Lutchmaya, Baron-Cohen, & Raggatt, 2002b). Geschwind's theory was that foetal testosterone might accelerate the growth of the right hemisphere at the expense of the left, which is usually dominant for language functions and which may also be of some significance for empathy.

Summary, and future work

Social neuroscience is now an important part of cognitive neuroscience. Studies of autism have contributed to this new field, and the literature reviewed earlier hints at the validity of an amygdala theory of autism. Future studies will be needed to test this more extensively. Secondly, future research will need to specify in greater detail which of the 13 nuclei in the amygdala are intact in autism, and which are impaired. Finally, the intriguing possibility that foetal testosterone mediates empathy through testosterone receptors in the amygdala and other parts of the 'social brain' will be an important hypothesis to test, when methods become available.

References:

- A.P.A. (1994). *DSM-IV Diagnostic and Statistical Manual of Mental Disorders, 4th Edition*. Washington DC: American Psychiatric Association.
- Abell, F, Krams, M, Ashburner, J, Passingham, R, Friston, K, Frackowiak, R, Happe, F, Frith, C, & Frith, U. (1999). The neuroanatomy of autism: a voxel-based whole brain analysis of structural scans. *Cognitive Neuroscience, 10*, 1647-1651.
- Adolphs, R, Sears, L, & Piven, J. (2001). Abnormal processing of social information from faces in autism. *Journal of Cognitive Neuroscience, 13*, 232-240.
- Adolphs, R, Tranel, D, Damasio, H, & Damasio, A. (1994). Impaired recognition of emotion in facial expressions following bilateral damage to the human amygdala. *Nature, 372*(669-672).
- Arnold, A P, & Gorski, R A. (1984). Gonadal steroid induction of structural sex differences in the CNS. *Annual Review of Neurosciences, 7*, 413-442.
- Asperger, H. (1944). Die "Autistischen Psychopathen" im Kindesalter. *Archiv fur Psychiatrie und Nervenkrankheiten, 117*, 76-136.
- Astington, J, Harris, P, & Olson, D. (1988). *Developing theories of mind*. New York: Cambridge University Press.
- Avis, J, & Harris, P. (1991). Belief-desire reasoning among Baka children: evidence for a universal conception of mind. *Child Development, 62*, 460-467.
- Bachevalier, J. (1991). An animal model for childhood autism: memory loss and socio-emotional disturbances following neonatal damage to the limbic system in monkeys. In C. Tamminga & S. Schulz (Eds.), *Advances in Neuropsychiatry and Psychopharmacology: Volume 1. Schizophrenia Research*. New York: Raven Press.
- Bachevalier, J. (1994). Medial temporal lobe structures and autism: a review of clinical and experimental findings. *Neuropsychologia, 32*, 627-648.
- Baron-Cohen, Jolliffe, T, Mortimore, C, & Robertson, M. (1997). Another advanced test of theory of mind: evidence from very high functioning adults with autism or Asperger Syndrome. *Journal of Child Psychology and Psychiatry, 38*, 813-822.
- Baron-Cohen, S. (1987). Perception in autistic children. In D. Cohen (Ed.), *Handbook of autism and pervasive developmental disorders*. New York: Wiley & Sons.
- Baron-Cohen, S. (1989c). Are autistic children behaviourists? An examination of their mental-physical and appearance-reality distinctions. *Journal of Autism and Developmental Disorders, 19*, 579-600.
- Baron-Cohen, S. (1989b). The autistic child's theory of mind: a case of specific developmental delay. *Journal of Child Psychology and Psychiatry, 30*, 285-298.
- Baron-Cohen, S. (1989a). Joint attention deficits in autism: towards a cognitive analysis. *Development and Psychopathology, 1*, 185-189.
- Baron-Cohen, S. (1991). Do people with autism understand what causes emotion? *Child Development, 62*, 385-395.
- Baron-Cohen, S. (1994). How to build a baby that can read minds: Cognitive mechanisms in mindreading. *Cahiers de Psychologie Cognitive/ Current Psychology of Cognition, 13*, 513-552.
- Baron-Cohen, S. (1995). *Mindblindness: an essay on autism and theory of mind*. Boston: MIT Press/Bradford Books.

- Baron-Cohen, S. (2002). The extreme male brain theory of autism. *Trends in Cognitive Sciences*, 6, 248-254.
- Baron-Cohen, S, & Goodhart, F. (1994). The Mindreading System: new directions for research. *Current Psychology of Cognition*, 13, 724-750.
- Baron-Cohen, S, & Goodhart, F. (1994). The "seeing leads to knowing" deficit in autism: the Pratt and Bryant probe. *British Journal of Developmental Psychology*, 12, 397-402.
- Baron-Cohen, S, & Hammer, J. (1997). Is autism an extreme form of the male brain? *Advances in Infancy Research*, 11, 193-217.
- Baron-Cohen, S, Leslie, A M, & Frith, U. (1985). Does the autistic child have a 'theory of mind'? *Cognition*, 21, 37-46.
- Baron-Cohen, S, O'Riordan, M, Jones, R, Stone, V, & Plaisted, K. (1999). A new test of social sensitivity: Detection of faux pas in normal children and children with Asperger syndrome. *Journal of Autism and Developmental Disorders*, 29, 407-418.
- Baron-Cohen, S, Richler, J, Bisarya, D, Gurunathan, N, & Wheelwright, S. (2003). The Systemising Quotient (SQ) : An investigation of adults with Asperger Syndrome or High Functioning Autism and normal sex differences. *Philosophical Transactions of the Royal Society, Series B, Special issue on "Autism : Mind and Brain"*, 358, 361-374.
- Baron-Cohen, S, & Ring, H. (1994). A model of the mindreading system: neuropsychological and neurobiological perspectives. In P. Mitchell & C. Lewis (Eds.), *Origins of an understanding of mind*. Hove, E.Sussex: Lawrence Erlbaum Associates.
- Baron-Cohen, S, Ring, H, Bullmore, E, Wheelwright, S, Ashwin, C, & Williams, S. (2000). The amygdala theory of autism. *Neuroscience and Behavioural Reviews*, 24, 355-364.
- Baron-Cohen, S, Ring, H, Moriarty, J, Shmitz, P, Costa, D, & Ell, P. (1994). Recognition of mental state terms: a clinical study of autism, and a functional neuroimaging study of normal adults. *British Journal of Psychiatry*, 165, 640-649.
- Baron-Cohen, S, Ring, H, Wheelwright, S, Bullmore, E, T., Brammer, M, J., Simmons, A, & Williams, S. (1999). Social intelligence in the normal and autistic brain: an fMRI study. *European Journal of Neuroscience*, 11, 1891-1898.
- Baron-Cohen, S, Tager-Flusberg, H, & Cohen, D (Eds.). (1993). *Understanding other minds: perspectives from autism*. Oxford: Oxford University Press.
- Baron-Cohen, S, & Wheelwright, S. (2004). The Empathy Quotient (EQ). An investigation of adults with Asperger Syndrome or High Functioning Autism, and normal sex differences. *Journal of Autism and Developmental Disorders*, 34, 163-175.
- Baron-Cohen, S, Wheelwright, S, Hill, J, Raste, Y, & Plumb, I. (2001). The 'Reading the Mind in the eyes' test revised version: A study with normal adults, and adults with Asperger Syndrome or High-Functioning autism. *Journal of Child Psychiatry and Psychology*, 42, 241-252.
- Baron-Cohen, S, Wheelwright, S, & Jolliffe, T. (1997). Is there a "language of the eyes"? Evidence from normal adults and adults with autism or Asperger syndrome. *Visual Cognition*, 4, 311-331.

- Baron-Cohen, S, Wheelwright, S, Stone, V, & Rutherford, M. (1999). A mathematician, a physicist, and a computer scientist with Asperger Syndrome: performance on folk psychology and folk physics test. *Neurocase*, 5, 475-483.
- Bates, E, Benigni, L, Bretherton, I, Camaioni, L, & Volterra, V. (1979). Cognition and communication from 9-13 months: correlational findings. In E. Bates (Ed.), *The emergence of symbols: cognition and communication in infancy*. New York: Academic Press.
- Bauman, M, & Kemper, T. (1994). *The Neurobiology of Autism*. Baltimore: Johns Hopkins.
- Bauman, M, & Kempner, T. (1985). Histoanatomic observation of the brain in early infantile autism. *Neurology*, 35, 866-874.
- Bauman, M, & Kempner, T. (1988). Limbic and cerebellar abnormalities: consistent findings in infantile autism. *Journal of Neuropathology and Experimental Neurology*, 47, 369.
- Bauman, M L, & Kemper, T L. (1994). Neuroanatomic observations of the brain in autism. In M. L. Bauman & T. L. Kemper (Eds.), *The Neurobiology of Autism* (pp. 119-145). Baltimore: John Hopkins University Press.
- Bauman, M L, & Kempner, T L. (1986). Developmental cerebellar abnormalities: A consistent finding in early infantile autism. *Neurology*, 36, 190.
- Bishop, D V M. (1993). Annotation: Autism, executive functions, and theory of mind: A neuropsychological perspective. *Journal of Child Psychology and Psychiatry*, 54, 279-293.
- Bolton, P, & Griffiths, P. (1997). Association of tuberous sclerosis of temporal lobes with autism and atypical autism. *The Lancet*, 349, 392-395.
- Bonda, E, Petrides, M, Ostry, D, & Evans, A. (1996). Specific involvement of human parietal systems and the amygdala in the perception of biological motion. *Journal of Neuroscience*, 15(3737-3744).
- Bowlby, J. (1969). *Attachment*. London: The Hogarth Press.
- Brentano, F. (1970). *Psychology from an empirical standpoint*. London: Routledge, and Kegan Paul.
- Brothers, L. (1990). The social brain: a project for integrating primate behaviour and neurophysiology in a new domain. *Concepts in Neuroscience*, 1, 27-51.
- Bruner, J. (1983). *Child's talk: learning to use language*. Oxford: Oxford University Press.
- Butterworth, G. (1991). The ontogeny and phylogeny of joint visual attention. In A. Whiten (Ed.), *Natural theories of mind*. Oxford: Blackwell.
- Collaer, M, & Hines, M. (1995). Human behavioural sex differences: A role for gonadal hormones during early development? *Psychological Bulletin*, 118, 55-107.
- Connellan, J, Baron-Cohen, S, Wheelwright, S, Ba'tki, A, & Ahluwalia, J. (2001). Sex differences in human neonatal social perception. *Infant Behavior and Development*, 23, 113-118.
- Courchesne, E, Townsend, J, Akshoomoff, N A, Yeung-Courchesne, R, Lincoln, A, J., Press, G, Murakami, J, James, H, Saitoh, O, Egaas, B, Haas, R, H., & Schreibman, L. (1994). A new finding: impairment in shifting attention in autistic and cerebellar patients. In S. H. Broman & J. Grafman (Eds.), *Atypical Cognitive Deficits in Developmental Disorders : Implications for Brain Function*. Hillsdale, New Jersey: Lawrence Erlbaum.

- Csibra, G, Gergely, G, Biro, S, Koos, O, & Brockbanck, M. (1999). Goal attribution without agency cues: the perception of 'pure reason' in infancy. *Cognition*, 72, 253-284.
- Damasio, A, Tranel, D, & Damasio, H. (1990). Individuals with sociopathic behaviour caused by frontal lobe damage fail to respond autonomically to socially charged stimuli. *Behavioural Brain Research*, 14, 81-94.
- De Long, G R. (1992). Autism, amnesia, hippocampus, and learning. *Neuroscience Behaviour Review*, 16, 63-70.
- Dennett, D. (1987). *The Intentional Stance*. Cambridge, Mass: MIT Press/Bradford Books.
- Ekman, P. (1992). Facial expressions of emotion: an old controversy and new findings. *Philos Trans R Soc Lond B Biol Sci*, 335, 63-69.
- Filipek, P A. (1999). Neuroimaging in the developmental disorders: the state of the science. *Journal of Child Psychology and Psychiatry*, 40(113 - 128).
- Fodor, J. (1983). *The modularity of mind*. Cambridge, USA: MIT/Bradford Books.
- Frith, U. (1991). *Autism and Asperger's Syndrome*. Cambridge: Cambridge University Press.
- Gelman, S, & Hirschfield, L. (1994). *Mapping the Mind*. Cambridge: Press Syndicate, University of Cambridge.
- Gergely, G, Nadasdy, Z, Gergely, C, & Biro, S. (1995). Taking the intentional stance at 12 months of age. *Cognition*, 56, 165-193.
- Geschwind, N, & Galaburda, A. (1987). *Cerebral lateralization*. Cambridge, MA: MIT Press.
- Geschwind, N, & Galaburda, A M. (1985). Cerebral lateralization. biological mechanisms, associations, and pathology: I. A hypothesis and a program for research. *Archive of Neurology*, 42, 428-459.
- Geschwind, N, & Galaburda, A M. (1985). Cerebral lateralization: biological mechanisms, associations and pathology. II. A hypothesis and a program for research. *Archives of Neurology*, 42, 521-552.
- Geschwind, N, & Galaburda, A M. (1985). Cerebral lateralization: biological mechanisms, associations and pathology. III. A hypotehsis and a program for research. *Archives of Neurology*, 42, 634-654.
- Gillberg, I, Bjure, J, Uvebrant, P, Vestergren, E, & Gillberg, C. (1993). SPECT in 31 children and adolescents with autism and autistic like syndromes. *European Child and Adolescent Psychiatry*, 2, 50-59.
- Goleman, D. (1995). *Emotional Intelligence*. New York: Bantam Books.
- Gross, C, Rocha-Miranda, C, & Bender, D. (1972). Visual properties of neurons in the inferotemporal cortex of the macaque. *Journal of Neurophysiology*, 35, 96-111.
- Happe, F. (1993). Communicative competence and theory of mind in autism: A test of Relevance Theory. *Cognition*, 48, 101-119.
- Happe, F, Ehlers, S, Fletcher, P, Frith, U, Johansson, M, Gillberg, C, Dolan, R, Frackowiak, R, & Frith, C. (1996). Theory of mind in the brain. Evidence from a PET scan study of Asperger Syndrome. *NeuroReport*, 8, 197-201.
- Harris, G W, & Levine, S. (1962). Sexual differentiation of the brain and its experimental control. *Journal of Physiology*, 181, 379-400.
- Harris, P, Johnson, C N, Hutton, D, Andrews, G, & Cooke, T. (1989). Young children's theory of mind and emotion. *Cognition and Emotion*, 3, 379-400.

- Hermelin, B., & O'Connor, N. (1970). *Psychological experiments with autistic children*. London: Pergamon Press.
- Hetzler, B., & Griffin, J. (1981). Infantile autism and the temporal lobe of the brain. *Journal of Autism and Developmental Disorders*, 9, 153-157.
- Hines, M., & Shipley, C. (1984). Prenatal exposure to Diethylstilbestrol (DES) and the development of sexually dimorphic cognitive abilities and cerebral lateralisation. *Developmental Psychology*, 20, 81-94.
- Hobson, R P. (1986). The autistic child's appraisal of expressions of emotion. *Journal of Child Psychology and Psychiatry*, 27, 321-342.
- Hood, B., Willen, J., & Driver, J. (1997). An eye-direction detector triggers shifts of visual attention in human infants. *Unpublished ms.*, Harvard University.
- Hyde, J S., & Linn, M C. (1988). Gender differences in verbal ability: A meta-analysis. *Psychological Bulletin*, 104, 53-69.
- I.C.D-10. (1994). *International classification of diseases* (10th ed.). Geneva, Switzerland: World Health Organisation.
- Johnson, S. (2000). The recognition of mentalistic agents in infancy. *Trends in Cognitive Sciences*, 4, 22-28.
- Kanner, L. (1943). Autistic disturbance of affective contact. *Nervous Child*, 2, 217-250.
- Karmiloff-Smith, A., Grant, J., Bellugi, U., & Baron-Cohen, S. (1995). Is there a social module? Language, face-processing and theory of mind in William's Syndrome and autism. *Journal of Cognitive Neuroscience*, 7, 196-208.
- Kawashima, R., Sugiura, M., Kato, T., Nakamura, A., Hatano, K., Ito, K., Fukuda, H., Kojima, s., & Nakamura, K. (1999). The human amygdala plays an important role in gaze monitoring. *Brain*, 122, 779-783.
- Kimura, D. (1999). *Sex and Cognition*. Cambridge, MA: MIT Press.
- Kling, A., & Brothers, L. (1992). The amygdala and social behavior. In J. Aggleton (Ed.), *Neurobiological aspects of emotion, memory, and mental dysfunction*. New York: Wiley-Liss, Inc.
- Leslie, A. (1995). ToMM, ToBy, and Agency: core architecture and domain specificity. In L. Hirschfeld & S. Gelman (Eds.), *Domain specificity in cognition and culture*. New York: Cambridge University Press.
- Leslie, A., & Keeble, S. (1987). Do six-month old infants perceive causality? *Cognition*, 25, 265-288.
- Leslie, A M. (1987). Pretence and representation: the origins of "theory of mind". *Psychological Review*, 94, 412-426.
- Leslie, A M., & Frith, U. (1988). Autistic children's understanding of seeing, knowing, and believing. *British Journal of Developmental Psychology*, 6, 315-324.
- Lutchmaya, S, Baron-Cohen, S, & Raggatt, P. (2002a). Foetal testosterone and eye contact in 12 month old infants. *Infant Behavior and Development*, 25, 327-335.
- Lutchmaya, S, Baron-Cohen, S, & Raggatt, P. (2002b). Foetal testosterone and vocabulary size in 18- and 24-month-old infants. *Infant Behavior and Development*, 24(4), 418-424.
- Maccoby, E., & Jacklin, N. (1974). *The Psychology of Sex Differences*. Stanford: Stanford University Press.
- Morris, J, Frith, C, Perrett, D, Rowland, D, Young, A, Calder, A, & Dolan, R. (1996). A differential neural response in the human amygdala to fearful and happy facial expressions. *Nature*, 383(812-815).

- Mundy, P, & Crowson, M. (1997). Joint attention and early social communication. *Journal of Autism and Developmental Disorders*, 27, 653-676.
- Ozonoff, S, Pennington, B, & Rogers, S J. (1990). Are there emotion perception deficits in young autistic children? *Journal of Child Psychology and Psychiatry*, 31, 343-363.
- Perner, J, & Wimmer, H. (1985). "John thinks that Mary thinks that..." Attribution of second-order beliefs by 5-10 year old children. *Journal of Experimental Child Psychology*, 39, 437-471.
- Pratt, C, & Bryant, P. (1990). Young children understand that looking leads to knowing (so long as they are looking into a single barrel). *Child Development*, 61, 973-983.
- Premack, D. (1990). The infant's theory of self-propelled objects. *Cognition*, 36, 1-16.
- Premack, D, & Woodruff, G. (1978). Does the chimpanzee have a 'theory of mind'? *Behaviour and Brain Sciences*, 4, 515-526.
- Rapin, I, & Katzman, R. (1998). Neurobiology of Autism. *Annals of Neurology*, 43, 7-14.
- Raven, J C. (1956). *Coloured Progressive Matrices*. London: H.K. Lewis and Co.
- Raymond, G, Bauman, M, & Kemper, T. (1996). Hippocampus in autism: a Golgi analysis. *Acta Neuropathol*, 91, 117-119.
- Rimland, B. (1964). *Infantile Autism : the syndrome and its implications for a neural theory of behaviour*. New York: Appleton-Century-Crofts.
- Rochat, P, Morgan, R, & Carpenter, M. (1997). Young infants' sensitivity to movement information specifying social causality. *Cognitive Development*, 12, 537-561.
- Rutter, M. (1978). Language disorder and infantile autism. In M. Rutter & E. Schopler (Eds.), *Autism: a reappraisal of concepts and treatment*. New York: Plenum.
- Sacks, O. (1994). *An anthropologist on Mars*. New York: Knopf.
- Scaife, M, & Bruner, J. (1975). The capacity for joint visual attention in the infant. *Nature*, 253, 265-266.
- Shantz, C. (1983). Social cognition. In P. Mussen (Ed.), *Handbook of Child Psychology: Vol 3: Cognitive Development* (pp. 495-555). New York: John Wiley and Sons.
- Tager-Flusberg, H. (1993). What language reveals about the understanding of minds in children with autism. In S. Baron-Cohen, H. Tager-Flusberg & D. Cohen, J, (Eds.), *Understanding other minds : perspectives from autism*. Oxford: Oxford University Press.
- Tager-Flusberg, H, Boshart, J, & Baron-Cohen, S. (1998). Reading the windows of the soul: evidence of domain specificity sparing in Williams syndrome. *Journal of Cognitive Neuroscience*, 10, 631-639.
- Tomasello, M. (1988). The role of joint-attentional processes in early language acquisition. *Language Sciences*, 10, 69-88.
- Tranel, D, & Hyman, B, T. (1990). Neuropsychological correlates of bilateral amygdala damage. *Arch Neurol*, 47, 349-355.
- Walker, A S. (1982). Intermodal perception of expressive behaviours by human infants. *Journal of Experimental Child Psychology*, 33, 514-535.
- Wechsler, D. (1939). *The Measurement of Adult Intelligence*. Baltimore: Williams and Wilkins.
- Wellman, H. (1990). *Children's theories of mind*. Cambridge, USA: Bradford/MIT Press.

- Wellman, H, & Bartsch, K. (1988). Young children's reasoning about beliefs. *Cognition*, 30, 239-277.
- Whalen, P J, Rauch, S L, Etcoff, N L, McInerney, S C, Lee, M B, & Jenike, M A. (1998). Masked presentations of emotional facial expressions modulate amygdala activity without explicit knowledge. *The Journal of Neuroscience*, 18, 411-418.
- Whiten, A. (1991). *Natural theories of mind*. Oxford: Basil Blackwell.
- Whiten, A. (1997). The Machiavellian Mindreader. In A. Whiten & R. W. Byrne (Eds.), *Machiavellian Intelligence II: Evaluations and Extensions*. Cambridge: CUP.
- Wicker, B, Michel, F, Henaff, M, & Decety, J. (1998). Brain regions involved in the perception of gaze: a PET study. *Neuroimage*, 8, 221-227.
- Williams, C, Barnett, A, & Meck, W. (1990). Organisational effects of early gonadal secretions on sexual differentiation in spatial memory. *Behavioural Neuroscience*, 104, 84-97.
- Wilson, J D, Foster, D W, Kronenberg, H M, & Larsen, P R (Eds.). (1998). *Williams Textbook of Endocrinology*. Philadelphia, London, Toronto, Montreal, Sydney, Tokyo: W B Saunders Company.
- Wimmer, H, & Perner, J. (1983). Beliefs about beliefs: Representation and constraining function of wrong beliefs in young children's understanding of deception. *Cognition*, 13, 103-128.
- Wing, L. (1981). Asperger Syndrome: a clinical account. *Psychological Medicine*, 11, 115-130.
- Wing, L, & Gould, J. (1979). Severe impairments of social interaction and associated abnormalities in children: epidemiology and classification. *Journal of Autism and Developmental Disorders*, 9, 11-29.
- Yirmiya, N, Sigman, M, Kasari, C, & Mundy, P. (1992). Empathy and cognition in high functioning children with autism. *Child Development*, 63, 150-160
- Young, A, Hellawell, D, De Wal, C, & Johnson, M. (1996). Facial expression processing after amygdectomy. *Neuropsychologia*, 34, 31-39.